Chapter 8 FRQ Classwork

1.

A population is modeled by a function P that satisfies the logistic differential equation

$$\frac{dP}{dt} = \frac{P}{5} \left(1 - \frac{P}{12} \right).$$

- (a) If P(0) = 3, what is $\lim_{t \to \infty} P(t)$? If P(0) = 20, what is $\lim_{t \to \infty} P(t)$?
- (b) If P(0) = 3, for what value of P is the population growing the fastest?
- (c) A different population is modeled by a function Y that satisfies the separable differential equation

$$\frac{dY}{dt} = \frac{Y}{5} \left(1 - \frac{t}{12} \right).$$

Find Y(t) if Y(0) = 3.

(d) For the function Y found in part (c), what is $\lim_{t \to \infty} Y(t)$?

2.

Consider the function $f(x) = \frac{1}{x^2 - kx}$, where k is a nonzero constant. The derivative of f is given by

$$f'(x) = \frac{k - 2x}{\left(x^2 - kx\right)^2}.$$

- (a) Let k = 3, so that $f(x) = \frac{1}{x^2 3x}$. Write an equation for the line tangent to the graph of f at the point whose x-coordinate is 4.
- (b) Let k = 4, so that $f(x) = \frac{1}{x^2 4x}$. Determine whether f has a relative minimum, a relative maximum, or neither at x = 2. Justify your answer.
- (c) Find the value of k for which f has a critical point at x = -5.
- (d) Let k = 6, so that $f(x) = \frac{1}{x^2 6x}$. Find the partial fraction decomposition for the function f. Find $\int f(x) dx$.

3.

Consider the differential equation $\frac{dy}{dx} = y^2(2x + 2)$. Let y = f(x) be the particular solution to the differential equation with initial condition f(0) = -1.

- (a) Find $\lim_{x\to 0} \frac{f(x)+1}{\sin x}$. Show the work that leads to your answer.
- (b) Use Euler's method, starting at x = 0 with two steps of equal size, to approximate $f\left(\frac{1}{2}\right)$.
- (c) Find y = f(x), the particular solution to the differential equation with initial condition f(0) = -1.

4.

Let f be the function defined for x > 0, with f(e) = 2 and f', the first derivative of f, given by $f'(x) = x^2 \ln x$.

- (a) Write an equation for the line tangent to the graph of f at the point (e, 2).
- (b) Is the graph of f concave up or concave down on the interval 1 < x < 3? Give a reason for your answer.
- (c) Use antidifferentiation to find f(x).