

## Chapter 8

1.

Consider the differential equation  $\frac{dy}{dx} = x^2 - \frac{1}{2}y$ .

(a) Find  $\frac{d^2y}{dx^2}$  in terms of  $x$  and  $y$ .

(b) Let  $y = f(x)$  be the particular solution to the given differential equation whose graph passes through the point  $(-2, 8)$ . Does the graph of  $f$  have a relative minimum, a relative maximum, or neither at the point  $(-2, 8)$ ? Justify your answer.

(c) Let  $y = g(x)$  be the particular solution to the given differential equation with  $g(-1) = 2$ . Find

$$\lim_{x \rightarrow -1} \left( \frac{g(x) - 2}{3(x+1)^2} \right).$$
 Show the work that leads to your answer.

(d) Let  $y = h(x)$  be the particular solution to the given differential equation with  $h(0) = 2$ . Use Euler's method, starting at  $x = 0$  with two steps of equal size, to approximate  $h(1)$ .

2.


Let  $f(x) = e^{2x}$ . Let  $R$  be the region in the first quadrant bounded by the graph of  $f$ , the coordinate axes, and the vertical line  $x = k$ , where  $k > 0$ . The region  $R$  is shown in the figure above.

(a) Write, but do not evaluate, an expression involving an integral that gives the perimeter of  $R$  in terms of  $k$ .

(b) The region  $R$  is rotated about the  $x$ -axis to form a solid. Find the volume,  $V$ , of the solid in terms of  $k$ .

(c) The volume  $V$ , found in part (b), changes as  $k$  changes. If  $\frac{dk}{dt} = \frac{1}{3}$ ,

determine  $\frac{dV}{dt}$  when  $k = \frac{1}{2}$ .



3.

Let  $f$  and  $g$  be the functions defined by  $f(x) = \frac{1}{x}$  and  $g(x) = \frac{4x}{1+4x^2}$ , for all  $x > 0$ .

(a) Find the absolute maximum value of  $g$  on the open interval  $(0, \infty)$  if the maximum exists. Find the absolute minimum value of  $g$  on the open interval  $(0, \infty)$  if the minimum exists. Justify your answers.

(b) Find the area of the unbounded region in the first quadrant to the right of the vertical line  $x = 1$ , below the graph of  $f$ , and above the graph of  $g$ .

## Chapter 9

4.

The function  $f$  has a Taylor series about  $x = 1$  that converges to  $f(x)$  for all  $x$  in the interval of convergence.

It is known that  $f(1) = 1$ ,  $f'(1) = -\frac{1}{2}$ , and the  $n$ th derivative of  $f$  at  $x = 1$  is given by

$$f^{(n)}(1) = (-1)^n \frac{(n-1)!}{2^n} \text{ for } n \geq 2.$$

- Write the first four nonzero terms and the general term of the Taylor series for  $f$  about  $x = 1$ .
- The Taylor series for  $f$  about  $x = 1$  has a radius of convergence of 2. Find the interval of convergence. Show the work that leads to your answer.
- The Taylor series for  $f$  about  $x = 1$  can be used to represent  $f(1.2)$  as an alternating series. Use the first three nonzero terms of the alternating series to approximate  $f(1.2)$ .
- Show that the approximation found in part (c) is within 0.001 of the exact value of  $f(1.2)$ .

5.

The function  $g$  has derivatives of all orders, and the Maclaurin series for  $g$  is


$$\sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+3} = \frac{x}{3} - \frac{x^3}{5} + \frac{x^5}{7} - \dots$$

- Using the ratio test, determine the interval of convergence of the Maclaurin series for  $g$ .
- The Maclaurin series for  $g$  evaluated at  $x = \frac{1}{2}$  is an alternating series whose terms decrease in absolute value to 0. The approximation for  $g\left(\frac{1}{2}\right)$  using the first two nonzero terms of this series is  $\frac{17}{120}$ . Show that this approximation differs from  $g\left(\frac{1}{2}\right)$  by less than  $\frac{1}{200}$ .
- Write the first three nonzero terms and the general term of the Maclaurin series for  $g'(x)$ .

6.

Let  $f(x) = \sin(x^2) + \cos x$ . The graph of  $y = |f^{(5)}(x)|$  is shown above.


- Write the first four nonzero terms of the Taylor series for  $\sin x$  about  $x = 0$ , and write the first four nonzero terms of the Taylor series for  $\sin(x^2)$  about  $x = 0$ .
- Write the first four nonzero terms of the Taylor series for  $\cos x$  about  $x = 0$ . Use this series and the series for  $\sin(x^2)$ , found in part (a), to write the first four nonzero terms of the Taylor series for  $f$  about  $x = 0$ .
- Find the value of  $f^{(6)}(0)$ .
- Let  $P_4(x)$  be the fourth-degree Taylor polynomial for  $f$  about  $x = 0$ . Using information from the graph of  $y = |f^{(5)}(x)|$  shown above, show that  $\left|P_4\left(\frac{1}{4}\right) - f\left(\frac{1}{4}\right)\right| < \frac{1}{3000}$ .



Graph of  $y = |f^{(5)}(x)|$

Chapter 10

7.




At time  $t$ , the position of a particle moving in the  $xy$ -plane is given by the parametric functions  $(x(t), y(t))$ , where  $\frac{dx}{dt} = t^2 + \sin(3t^2)$ . The graph of  $y$ , consisting of three line segments, is shown in the figure above. At  $t = 0$ , the particle is at position  $(5, 1)$ .

- Find the position of the particle at  $t = 3$ .
- Find the slope of the line tangent to the path of the particle at  $t = 3$ .
- Find the speed of the particle at  $t = 3$ .
- Find the total distance traveled by the particle from  $t = 0$  to  $t = 2$ .

8.

The graphs of the polar curves  $r = 3$  and  $r = 3 - 2\sin(2\theta)$  are shown in the figure above for  $0 \leq \theta \leq \pi$ .



- Let  $R$  be the shaded region that is inside the graph of  $r = 3$  and inside the graph of  $r = 3 - 2\sin(2\theta)$ . Find the area of  $R$ .
- For the curve  $r = 3 - 2\sin(2\theta)$ , find the value of  $\frac{dx}{d\theta}$  at  $\theta = \frac{\pi}{6}$ .
- The distance between the two curves changes for  $0 < \theta < \frac{\pi}{2}$ .

Find the rate at which the distance between the two curves is changing with respect to  $\theta$  when  $\theta = \frac{\pi}{3}$ .

- A particle is moving along the curve  $r = 3 - 2\sin(2\theta)$  so that  $\frac{d\theta}{dt} = 3$  for all times  $t \geq 0$ . Find the value of  $\frac{dr}{dt}$  at  $\theta = \frac{\pi}{6}$ .

9.

At time  $t$ , a particle moving in the  $xy$ -plane is at position  $(x(t), y(t))$ , where  $x(t)$  and  $y(t)$  are not explicitly given. For  $t \geq 0$ ,  $\frac{dx}{dt} = 4t + 1$  and  $\frac{dy}{dt} = \sin(t^2)$ . At time  $t = 0$ ,  $x(0) = 0$  and  $y(0) = -4$ .

- Find the speed of the particle at time  $t = 3$ , and find the acceleration vector of the particle at time  $t = 3$ .
- Find the slope of the line tangent to the path of the particle at time  $t = 3$ .
- Find the position of the particle at time  $t = 3$ .
- Find the total distance traveled by the particle over the time interval  $0 \leq t \leq 3$ .